

Cole Crops Program 2008

Including Broccoli, Cauliflower, & Cabbage

1. LTB products must be applied according to the procedures outlined below:
2. After the spray tank is mixed with other chemicals and/or water, **the pH must be neutral or slightly acidic (pH 7.0 or less)**. LTB products are to be added to tank lastly.
3. If rainfall sufficient to wash off product occurs within 4 hours of application, **re-treatment is required**.
4. Application timing is very important, so **the earlier application is strongly recommended**.

Product Used	Application Timing	Minimum Application Rate	Benefits (as compared to untreated check)
MegaGro	Foliar spray the flats just before transplanting OR in "drench" water at transplanting	Dilution ratio: 13 oz. / 100 gal. water	<ul style="list-style-type: none"> • Improves vigor of transplant • Promotes better root development
	2 true leaves if seeded; or when new growth starts after transplanted	2 oz. / acre as foliar spray**	
HappyGro + PhotoGro + BOMO	1) At bud formation 2) Additional sprays in 14-21 day intervals	2 oz. / acre** of each 5-10 oz. / acre	<ul style="list-style-type: none"> • Enhances size and quality
Weather King	21-28 days after MegaGro	2 oz / acre**	<ul style="list-style-type: none"> • Increases tolerance to environmental stresses

**** Note: If application spray volume is greater than 15 gallons per acre, use the dilution rate of 13 oz. per 100 gallons of water.**