

Fruiting Vegetables Program 2008

Including Tomatoes, Peppers, & Eggplant

1. LTB products must be applied according to the procedures outlined below:
2. After the spray tank is mixed with other chemicals and/or water, **the pH must be neutral or slightly acidic (pH 7.0 or less)**. LTB products are to be added to tank lastly.
3. If rainfall sufficient to wash off product occurs within 4 hours of application, **re-treatment is required**.
4. Application timing is very important, so **the earlier application is strongly recommended**.

Product Used	Application Timing	Minimum Application Rate	Benefits (as compared to untreated check)
MegaGro	1) Spray in flats just before transplanting OR in "drench" water at transplanting 2) Repeat in 7-10 days (4 leaves)	Dilution ratio: 13 oz / 100 gal. water	<ul style="list-style-type: none"> • Improves vigor of transplant • Promotes better root development
HappyGro + PhotoGro + BOMNZN	14-21 days after MegaGro and can be re-applied every 10 days until harvest	2 oz. / acre** of each 2-15 oz. / acre	<ul style="list-style-type: none"> • Early application initiates more flower buds • During bloom will set more fruit
Weather King	21-28 days after HappyGro	2 oz / acre **	<ul style="list-style-type: none"> • Increases tolerance to environmental stresses

**** Note:** If application spray volume is greater than 15 gallons per acre, use the dilution rate of 13 oz. per 100 gallons of water.