

Nut Tree Program 2008

Including Almond, Walnut, Pistachio, and Pecan

1. LTB products must be applied according to the procedures outlined below:
2. After the spray tank is mixed with other chemicals and/or water, **the pH must be neutral or slightly acidic (pH 7.0 or less)**. LTB products are to be added to tank lastly.
3. If rainfall sufficient to wash off product occurs within 4 hours of application, **re-treatment is required**.
4. Application timing is very important, so **the earlier application is strongly recommended**.

Product Used	Application Timing	Minimum Application Rate	Benefits (as compared to untreated check)
MegaGro	1 st : 30 days prior to bud break 2 nd : Nuts at 1/4 inch diameter (1/8 inch for pistachio)	13 oz./100 gallons	<ul style="list-style-type: none"> • Promotes new root development
HappyGro + PhotoGro	1 st : Bud swell 2 nd : 14 days after 1 st application	13 oz./100 gallons of each	<ul style="list-style-type: none"> • Promotes nut set • Promotes pollination
BOMO + OK Degree	1 st : Flowering 2 nd : Every 7-10 days after 1 st application	64 oz./ acre 13 oz./100 gallons	<ul style="list-style-type: none"> •

Note: In the off year of alternate bearing crops, to increase nut set, use HappyGro three times starting at flower bud every 7-10 days.