

Onion Program 2008

Including Garlic, Leek, (seed crop see note*)

1. LTB products must be applied according to the procedures outlined below:
2. After the spray tank is mixed with other chemicals and/or water, **the pH must be neutral or slightly acidic (pH 7.0 or less)**. LTB products are to be added to tank lastly.
3. If rainfall sufficient to wash off product occurs within 4 hours of application, **re-treatment is required**.
4. Application timing is very important, so **the earlier application is strongly recommended**.

Product Used	Application Timing	Minimum Application Rate	Benefits (as compared to untreated check)
MegaGro	2-4 inch sprout	2 oz. / acre as foliar spray**	<ul style="list-style-type: none"> • Improves root system • Enhances internal plant metabolism
PhotoGro + BOMO	2 weeks after MegaGro and spray every 2 weeks	2 oz / acre** 5-10 oz. / acre	<ul style="list-style-type: none"> • Enhances vigor & growth of bulb • Enhances photosynthesis and energy production • Increases bulb weight
Weather King	21-28 days after MegaGro application	2 oz. / acre**	<ul style="list-style-type: none"> • Increases tolerance to environmental stresses

*Note: In a seed crop, after MegaGro and PhotoGro, apply 4 oz of HappyGro in 25-30 gal water 7 days after full bloom.

** Note: If application spray volume is greater than 15 gallons per acre, use the dilution rate of 13 oz. per 100 gallons of water.