

Stone Fruit Program 2008

Including Peach, Plum, & Apricot

1. LTB products must be applied according to the procedures outlined below:
2. After the spray tank is mixed with other chemicals and/or water, **the pH must be neutral or slightly acidic (pH 7.0 or less)**. LTB products are to be added to tank lastly.
3. If rainfall sufficient to wash off product occurs within 4 hours of application, **re-treatment is required**.
4. Application timing is very important, so **the earlier application is strongly recommended**.

Product Used	Application Timing	Minimum Application Rate	Benefits (as compared to untreated check)
HappyGro* + PhotoGro + BOMNZN	At petal fall	13 oz./100 gallons of each 64 oz./acre	<ul style="list-style-type: none"> • Promotes fruit set • Promotes pollination
MegaGro	1 st : 30 days prior to bud break 2 nd : Small fruit set	13 oz./100 gallons	<ul style="list-style-type: none"> • Promotes new root development
BOMO + OK Degree	1 st : 7-14 days pre-bloom 2 nd : Full bloom 3 rd : Small fruit set	64 oz./ acre 13 oz./100 gallons	<ul style="list-style-type: none"> •

* Note: During off years of alternate bearing crops, to increase the fruit set, apply HappyGro twice-- first at swollen bud then petal fall.